complete list of medications that may cause photosensitivity

LUMAVANCE™ LED FACEMASK

DRUG TYPE	SPECIFIC GROUP OR COMMON NAME	COMMENTS
Anti-Arthritic	Gold 50 or Ridaura	If YES, the treatment cannot be administered
Anti-Arthritic or Immunosuppressant	Azathioprine (Imuran, Azasan)	If YES, the treatment can be administered as long as the medication has not been taken within the last 5 days.
Anti-Arrhythmic	Amiodarone (Cordarone, Pacerone), Aratac	If YES, the treatment can be administered as long as the medication has not been taken within the last 5 days.
	Quinidine	If YES, and the client is currently on the medication it is at the discretion of the client as to whether they commence the treatment. There is a 10/100 chance of a light reaction. If the client has stopped taking the medication for ≥ 5 days, then the treatment can be administered.
Antibiotics	Fluoroquinolones: Ciprofloxacin (Cipro), Levofloxacin (Levaquin), Lomefloxacin (Maxaquin),Norfloxacin (Noroxin), Ofloxacin (Floxin) Tetracyclines: Demeclocycline (Declomycin), Doxycycline (Vibramycin), Minocycline (Minocin), Oxytetracycline (Terramycin) Others: Azithromycin (Zithromax), Capreomycin (Capastat), Ceftazidime (Fortaz),cycloserine (Seromycin),Metronidazole (Flagyl), nalidixic acid (NegGram), pyrazinamide, sulfamethoxazole/trimethoprim (Bactrim)	If YES, the treatment can be administered as long as the medication has not been taken within the last 5 days.
Anti-Cancer	Bexarotene (Targretin), Capecitabine (Xeloda), Dacarbazine (DTIC), Epirubicin (Ellence), Fluorouracil (5-FU), Interferon alfa (Intron A, Alferon-N), Methotrexate (Mexate), Pentostatin (Nipent), Procarbazine (Matulane), Tretinoin, oral (Vesanoid), Vinblastine (Velban, Velbe)	If YES, and the client is currently on the medication it is at the discretion of the client as to whether they commence the treatment. There is between a 1 /100 and 5/100 chance of a light reaction. If the client has stopped taking the medication for ≥ 5 days, then the treatment can be administered.
Anticonvulsants	Carbamazepine (Tegretol), Felbamate (Felbatol), Gabapentin (Neurontin), Lamotrigine (Lamictal), Oxcarbazepine (Trileptal), Topiramate (Topamax), Valproic acid (Depakene)	If YES, and the client is currently on the medication it is at the discretion of the client as to whether they commence the treatment. There is a 1/100 chance of a light reaction. If the client has stopped taking the medication for ≥ 5 days, then the treatment can be administered.
Antifungals	Flucytosine (Ancobon), Griseofulvin (Fulvicin, Gris-PEG), Terconazole (Terazol) Voriconazole (VFEND)	If YES, the treatment can be administered as long as the medication has not been taken within the last 5 days.
Antihistamines	Cetirizine (Zyrtec), Diphenhydramine (Benadryl), Loratadine (Claritin), Promethazine (Phenergan)	If YES, the treatment can be administered as long as the medication has not been taken within the last 5 days.
Antihypertensives	Antihypertensives Captopril (Capoten), Diltiazem (Cardizem, Tiazac), Enalapril (Vasotec), Nifedipine (Procardia), Sotalol (Betapace)	If YES, the treatment can be administered as long as the medication has not been taken within the last 5 days.
Antimalarial	Chloroquine (Aralen), Hydroxychloroquine (Plaquenil), Pyrimethamine (Daraprim), Pyrimethamine/sulfadoxine (Fansidar), Quinine	If YES, the treatment can be administered as long as the medication has not been taken within the last 5 days
Antipsychotics	Phenothiazines: Chlorpromazine (Thorazine), Fluphenazine (Prolixin), Perphenazine (Trilafon), Prochlorperazine (Compazine), Thioridazine (Mellaril), Trifluoperazine (Stelazine)	If YES, and the client is currently on the medication it is at the discretion of the client as to whether they commence the treatment. There is between a 2/100 and 3/100 chance of a light reaction. If the client has stopped taking the medication for ≥ 5 days, then the treatment can be administered.

Antiretroviral	Ritonavir (Norvir), Saquinavir (Fortovase, Invirase), Zalcitabine (Hivid)	If YES, and the client is currently on the medication it is at the discretion of the client as to whether they commence the treatment. There is approximately a 2/100 chance of a light reaction. If the client has stopped taking the medication for ≥ 5 days, then the treatment can be administered.
Antiviral	Amantadine (Symmetrel), Acyclovir (Zovirax)	If YES, and the client is currently on the medication it is at the discretion of the client as to whether they commence the treatment. There is approximately a 1/100 chance of a light reaction. If the client has stopped taking the medication for ≥ 5 days, then the treatment can be administered.
Cardiovascular	Thiazide diuretics: Bendroflumethiazide (Corzide), Chlorthalidone (Thalitone), Hydrochlorothiazide (Microzide), Hydroflumethiazide (Diucardin), Indapamide (Lozol), Methyclothiazide (Enduron), Metolazone (Zaroxolyn), Polythiazide (Renese) Diuretics, Other: Furosemide (Lasix), Triamterene (Dyrenium)	If YES, the treatment can be administered as long as the medication has not been taken within the last 5 days.
Lipid regulators Other	Fenofibrate (Tricor)	If YES, and the client is currently on the medication it is at the discretion of the client as to whether they commence the treatment. There is a 10/100 chance of a light reaction. If the client has stopped taking the medication for ≥ 5days then the treatment can be administered.
Non-steroidal antiinflammatory (NSAIDs) Analgesics	Diclofenac (Voltaren, Cataflam), Naproxen (Anaprox)	If YES, and the client is currently on the medication it is at the discretion of the client as to whether they commence the treatment. There is a <1/100 chance of a light reaction. If the client has stopped taking the medication for ≥ 1 day, then the treatment can be administered.
Sedatives	Alprazolam (Xanax), Chlordiazepoxide (Librium), Zaleplon (Sonata), Zolpidem (Ambien)	If YES, and the client is currently on the medication it is at the discretion of the client as to whether they commence the treatment. There is a 1/100 chance of a light reaction. If the client has stopped taking the medication for ≥ 5 days, then the treatment can be administered.
Skin agents (acne)	Isotretinoin (Accutane, Roaccutane) Tretinoin topical (Renova, Retin-A) Tazarotene (Tazorac)	If YES, and the client is currently on the medication it is at the discretion of the client as to whether they commence the treatment. There is between a 5/100 and a 10/100 chance of a light reaction. If the client has stopped taking the medication for ≥ 5 days, then the treatment can be administered.
Skin agents (hair)	Coal tar, Minoxidil (Rogaine)	If YES, and the client is currently on the medication it is at the discretion of the client as to whether they commence the treatment. There is < 0.5/100 chance of a light reaction. If the client has stopped taking the medication for ≥ 5 days, then the treatment can be administered.